

RESEARCH

Open Access



Approach to fever in children among final-year nursing students: a multicenter survey

Gregorio P. Milani^{1,2*}, Antonio Corsello^{1,2}, Marta Fadda³, Ilaria Falvo³, Mario G. Bianchetti⁴, Diego Peroni⁵, Elena Chiappini⁶, Barbara Cantoni⁷, Patrizio Sannino⁷, Anne Destrebecq⁸ and Paola Marchisio^{1,9}

Abstract

Background Unfounded concerns regarding fever are increasingly observed among nurses worldwide. However, no study has so far explored the preferred approach towards pediatric fever among nursing students. Therefore, we aimed to investigate the attitude towards pediatric fever among final-year nursing students.

Methods Between February and June 2022, final-year nursing students of 5 Italian university hospitals were asked to answer an online survey on their approach to fever in children. Both quantitative and qualitative methods were utilized. Multiple regression models were employed to explore the existence of moderators on fever conceptions.

Results The survey was filled in by 121 nursing students (response rate 50%). Although most students (98%) do not consider discomfort to treat fever in children, only a minority would administer a second dose of the same antipyretic in nonresponsive cases (5.8%) or would alternate antipyretic drugs (13%). Most students would use physical methods to decrease fever (84%) and do not think that fever has mainly beneficial effects in children (72%). The own knowledge adequacy on fever was inversely associated (OR 0.33, 95% CI 0.13–0.81) with the beliefs that high fever might lead to brain damage. No further predictive variable was significantly associated with the concern that fever might be associated with brain damage, the advice of physical methods use, and the assumption that fever has mostly positive effects.

Discussion This study shows for the first time that misconceptions and inappropriate attitudes towards fever in children are common among final-year nursing students. Nursing students could potentially be ideal candidates for improving fever management within clinical practice and amongst caregivers.

Keywords Education, Children, Fever phobia, Inappropriate treatment, Evidence, Physical methods, Education

*Correspondence:

Gregorio P. Milani
milani.gregoriop@gmail.com

¹Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy

²Pediatric Unit, Fondazione IRCCS Ca'Granda Ospedale Maggiore Policlinico, via della Commenda 9, Milan 20122, Italy

³Institute of Public Health, Faculty of Biomedical Sciences, Università della Svizzera italiana, Lugano, Switzerland

⁴Faculty of Biomedical Sciences, Università della Svizzera Italiana, Lugano, Switzerland

⁵Department of Clinical and Experimental Medicine, Section of Pediatrics, University of Pisa, Pisa, Italy

⁶Pediatric Infectious Disease Unit, Department of Health Sciences, Meyer Children's University Hospital, University of Florence, Florence, Italy

⁷Direzione Professioni Sanitarie, Fondazione IRCCS Ca'Granda Ospedale Maggiore Policlinico, Milan, Italy

⁸Department of Biomedical Sciences for Health, University of Milan, Milan, Italy

⁹Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Fever in children is one of the most frequent complaints for care seeking [1]. It usually results from an infection and is a cardinal part of the inflammatory response [2]. As such, fever has a relevant role in fighting infection [3]. However, since defense mechanisms can go awry in a few cases, fever can be associated with an increased metabolic rate and distress, and affect the overall wellbeing of a child [4]. Therefore, the primary goal of fever treatment is to improve the child's comfort rather than modifying the body temperature [5].

Unfounded concerns and inappropriate attitudes towards fever have been increasingly observed among nurses [6], who play a key role both in the care of feverish children and in providing the information to caregivers [7]. For instance, several studies conducted among nurses have reported the use of physical antipyresis (such as sponges using cold water) or alternating antipyretic drugs to reduce body temperature [8–10]. These non-evidence-based behaviors, which mainly rely on the assumptions that fever is per se noxious, fail in the pursuit of decreasing body temperature and put children at risk of side effects (e.g., distress or drug overdose) [11, 12].

Despite educational interventions implemented to address this issue [10], a gap still exists between the available recommendations and attitudes towards fever among nurses [13, 14]. Nursing students might represent an ideal population for interventions aimed at transferring evidence-based recommendations into day-to-day clinical practice [13, 14].

However, no study has so far explored the preferred approach towards pediatric fever among nursing students. For this purpose, we carried out a survey to investigate the attitude towards pediatric fever among final-year nursing students.

Methods

Between February and June 2022, we conducted a cross-sectional study among final-year nursing students of 5 Italian university hospitals: Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico (Milan), Presidio Ospedaliero, ASST Rhodense (Rho), Istituto Nazionale dei Tumori (Milan), Ospedale Maggiore di Crema (Crema), and ASST Valle Olona (Busto Arsizio). Students were invited to answer a survey composed by both closed and open-ended questions regarding fever in otherwise healthy children.

A link to access an anonymous questionnaire on Google Form was sent to the institutional email of all eligible participants. Two further reminders were sent after 10 and 20 days, respectively.

To develop the questionnaire, we initially reviewed the seminal survey on child fever proposed by Barton D. Schmitt [15]. This survey has been utilized in over

70 studies globally [6]. Next, the survey was modified slightly by four international experts on child fever (M.F., M.G.B., E.C., and P.M.) and adapted for nursing students. The modified survey was then tested by six researchers, including two nurses, two board-certified pediatricians, and two science communicators, and two questions were slightly revised based on their feedback. Finally, two students completed the developed questionnaire independently on two occasions, with a 10-day interval between each administration. The intra-rater reproducibility was found to be 98%.

Briefly, the following information was collected by 15 close-ended questions: (1) age and gender; (2) training related to fever received during the nursing program and own know-how adequacy on fever; (3) attitude towards fever management in children. Six questions included Likert scale answers ("Very inadequate", "Inadequate", "Adequate" and "Very adequate", or "Very unlikely", "Unlikely", "Likely" and "Very likely", or "Strongly disagree", "Disagree", "Agree", "Strongly agree"). Moreover, open-ended questions explored second-hand or personal relevant experience related to child's fever. The full questionnaire is provided in the online supplementary material.

Statistical analysis

Data are reported as absolute frequency and percentage or as median and interquartile, as appropriate. Dichotomous variables were compared by Fisher exact test.

To investigate the existence of moderators on fever conceptions, the 4-digit scales of the answers related to (1) possible brain damage associated to fever, (2) the use of physical methods to decrease body temperature and (3) the beneficial effects of fever, were collapsed into 2-digit measures. Multiple logistic regression models were employed to assess the associations between the answers to the above-mentioned issues (dependent variables) and age, gender, teaching time on fever (<1 vs. ≥1 h), own know-how adequacy on fever ("very inadequate" + "inadequate" vs. "adequate" + "very adequate"), second-hand or personal relevant experience related to child's fever (independent variables). Odds ratios (ORs) with 95% confidence intervals (95% CIs) were calculated. A two-tailed $p < 0.05$ was assumed as significant. Statistical analyses were performed using "R" program (version 3.5.3, 2019).

Qualitative analysis

An inductive-deductive analysis of all open-ended answers was performed to define experiential patterns by two authors (MF and IF). The analysis comprised four phases: (1) data cleaning and data reorganization, during which we differentiated between personal and second-hand experiences; (2) inductive analysis, during which

Table 1 Characteristics of the 121 nursing students participating the study. Data are given as median and interquartile range or absolute frequency and percentage

| Age, years | 24 [22–25] |
|--|------------|
| Gender | |
| Males | 21 (17) |
| Females | 100 (83) |
| Teaching time dedicated to fever | |
| >5 h | 4 (3.3) |
| >2–5 h | 22 (18) |
| 1–2 h | 46 (38) |
| <1 h | 31 (26) |
| No time | 18 (15) |
| Topic fever deepened by means of a textbook, yes | 68 (56) |
| Own know-how adequacy on fever | |
| Very adequate | 11 (9.0) |
| Adequate | 58 (48) |
| Inadequate | 48 (40) |
| Very inadequate | 4 (3.3) |

Table 2 Management of fever in children by nursing students. Data are given as absolute frequency and percentage

| N | 121 |
|--|----------|
| Criterion for treating fever in children | |
| Body temperature | |
| >36.7 °C | 0 (0.0) |
| >37.4 °C | 26 (14) |
| >37.9 °C | 59 (47) |
| >38.4 °C | 28 (32) |
| >38.9 °C | 3 (4.9) |
| I would consider discomfort rather than the temperature | 6 (2.4) |
| Most prescribed antipyretic drug | |
| Acetaminophen | 110 (91) |
| Ibuprofen | 5 (4.1) |
| Salicylates | 6 (5.0) |
| Second dose of the same antipyretic in nonresponsive cases | |
| Very likely | 7 (5.8) |
| Likely | 0 (0.0) |
| Unlikely | 64 (53) |
| Very unlikely | 50 (41) |
| Alternating antipyretic regimen | |
| Very likely | 1 (0.8) |
| Likely | 15 (12) |
| Unlikely | 53 (44) |
| Very unlikely | 52 (43) |

we developed categories that represented frequent content patterns; (3) deductive analysis, where we reduced and re-labelled the categories to avoid redundancies and deductively coded all answers accordingly; and (4) thematic analysis in which we identified overarching themes and organized the results under four main types of experience. This approach is consistent with the thematic

approach described by Braun and Clarke [16, 17]. Further details are described elsewhere [18].

Ethical statement

The study was approved by the Institutional Ethics Committee of the University of Milan. Since no patient was involved, a single Ethics Committee approval was considered sufficient. All enrolled students provided a written informed consent to participate. No incentive or compensation was given to participants who full-filled the survey. The study was conducted according to the principles of Helsinki declaration and later amendments.

Results

A total of 241 final-year nursing students were invited to participate and 121 filled-in the survey (response rate 50%). Most participants were female (N=100, 82%). The median age of participants was 24 [interquartile range: 22–25] years. Many students evaluate their own know-how on fever (Table 1) as adequate or very adequate (N=69, 57%). Although most nursing students (N=115, 98%) do not consider discomfort rather than body temperature to treat fever in children (Table 2), only a minority would administer a second dose of the same antipyretic in nonresponsive cases (N=7, 5.8%) or would altern antipyretics (N=16, 13%). Most students do not think that fever can lead to brain damage (N=89, 74%). On the other hand, most participants would use physical methods to decrease fever (N=101, 83%) and do not think that fever has mainly beneficial effects in children (N=87, 72%), as shown in Table 3.

Nine and 15 respondents reported a relevant personal episode of fever from their childhood or about another child, respectively. The open answers pointed out that nursing students had relevant experiences associated to high body temperature, complications of fever (e.g., febrile seizures), etiology of fever, and management of high body temperature (details are provided in the supplementary online material). High body temperature and consequences of fever are the most frequently relevant experiences disclosed by participants.

The full results of the qualitative analyses (eTable 1) and of the logistic regression models (eTable 2–4) are reported within the online supplementary material. An adequate/very adequate own know-how on fever was inversely associated (OR 0.33, 95% CI 0.13–0.81) with the beliefs that high fever might lead to brain damage. No further predictive variable was significantly associated with the concern that high fever might lead to brain damage, the advice of physical methods and the assumption that fever has mostly positive effects.

Table 3 Answers of nursing students to questions on the assumption that fever can cause cerebral damage, on the use of physical methods to decrease the fever and on the belief that fever has mostly positive effects in children. Data are given as absolute frequency and percentage

| N | 121 |
|-------------------------------------|---------|
| Fever can cause cerebral damage | |
| Very likely | 6 (5.0) |
| Likely | 26 (21) |
| Unlikely | 80 (66) |
| Very unlikely | 9 (7.4) |
| Use of physical methods | |
| Very likely | 41 (34) |
| Likely | 60 (50) |
| Unlikely | 16 (13) |
| Very unlikely | 4 (3.3) |
| Fever has mostly beneficial effects | |
| Strongly agree | 1 (0.8) |
| Agree | 33 (27) |
| Disagree | 70 (58) |
| Strongly disagree | 17 (14) |

Discussion

This is the first study investigating the approach to pediatric fever among nursing students. The results indicate that misconceptions and improper approach towards fever in children are prevalent among final-year nursing students.

More than 40 years ago, Schmitt administered a questionnaire to caregivers of children visiting a hospital in Denver and noted unrealistic concerns about the possible consequences of fever. This attitude was also associated to a tendency to overtreat fever in most of the respondents [15]. Subsequent studies showed that this phenomenon, also known as “fever phobia”, was not limited to caregivers, but was common even among nurses in many countries [19–22]. This study reveals that merely 5% of nursing students view fever as a beneficial mechanism for children, with the majority instead recommending non-evidence-based interventions to reduce body temperature. These findings highlight the presence of fever phobia also within the nursing student population.

Several results from this survey are consistent with those of a previous study [18] we conducted among final-year medical students. Like medical students, a minority of nursing students consider discomfort, rather than body temperature, to be the primary indicator for treating fever (Fig. 1). However, more than 80% of both nursing and medical students would not administer an additional dose of antipyretic medication or switch to another antipyretic drug for a child with a persistently high body temperature. Furthermore, approximately 20% of nursing and medical students believe that fever can lead to brain damage, and more than half recommend physical methods to reduce body temperature. Finally, the lack of an

association between misconceptions and personal or second-hand experiences related to childhood fever in both groups of students supports the notion that individual learned traits have a negligible impact on the approach to fever [23]. Conversely, this study indicates that a minority of nursing students believe that fever is beneficial to children, whereas approximately half of the medical students (49%) agree with this evidence. These findings indirectly support the results of a recent systematic review, which suggests that misconceptions about fever are similarly prevalent among physicians and nurses, with the exception that nurses are less aware of the potential beneficial effects of fever [6].

The results of this study emphasize the potential benefits of training in the management of fever. A majority of nursing students (>70%) reported a limited amount of teaching time (less than 1 h or no time at all) devoted to child fever during their academic training. On the other hand, the only variable found to be associated with less frequent beliefs that fever may result in brain damage was the perception of an adequate knowledge on this subject. These data are relevant, given the crucial role that nurses play in educating parents about fever and managing it in various healthcare contexts, including outpatient clinics, emergency units, and pediatric wards [7]. If subsequent international surveys confirm the results of this study, then training on the management of child fever should receive greater attention in nursing academic programs worldwide.

This study has limitations. It was conducted exclusively in Italy and the sample size was limited. Furthermore, the responses provided by the students might not entirely represent the day-to-day clinical approach and




Fig. 1 Conceptions and attitudes of nursing and medical students [18]

might have been influenced by the desire to respond in a socially acceptable way. Finally, we could not investigate the possible role of regional differences on fever approach.

Conclusion

This study shows that misconceptions and inappropriate attitudes towards fever in children are common among final-year nursing students. This population might become an ideal candidate to improve fever management in the clinical practice and among caregivers, as the conceptions and behaviors of students play a crucial role in shaping their future work [24]. To this end, future research should address the impact of educational interventions on nursing practice going forward.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12912-023-01263-3>.

Supplementary Material 1

Acknowledgements

none.

Author Contribution

G.P.M.: study conception and design, analysis, data interpretation, writing of the first draft of the manuscript. A.C.: investigation, data collection and interpretation, prepared Fig. 1, review of the draft. M.F. and I.F.: study design, qualitative analysis, data interpretation, review of the draft. M.G.B.: study design, data interpretation, writing of the first draft of the manuscript. D.P., E.C., B.C., P.S., A.D. and P.M.: study design, data collection and interpretation, review of the draft. All authors gave the final approval of the version to be submitted.

Funding

The study was partially funded by the Italian Ministry of Health (Ricerca Corrente).

Data Availability

upon reasonable request to the corresponding author.

Declarations

Ethics approval and consent to participate

The study was approved by the Ethical Committee of the University of Milan and all participants gave their written consent to participate. The study was conducted according to the principles of Helsinki declaration and later amendments.

Consent for publication

not applicable.

Competing interests

The authors declare that they have no competing interests.

List of abbreviations

none.

Received: 10 November 2022 / Accepted: 22 March 2023

Published online: 13 April 2023

References

1. de Bont EG, Lepot JM, Hendrix DA, Loonen N, Guldemond-Hecker Y, Dinant GJ, Cals JW. Workload and management of childhood fever at general practice out-of-hours care: an observational cohort study. *BMJ Open*. 2015;5(5):e007365.
2. Evans SS, Repasky EA, Fisher DT. Fever and the thermal regulation of immunity: the immune system feels the heat. *Nat Rev Immunol*. 2015;15(6):335–49.
3. Rice P, Martin E, He JR, Frank M, DeTolla L, Hester L, O'Neill T, Manka C, Benjamin I, Nagarsekar A, et al. Febrile-range hyperthermia augments neutrophil accumulation and enhances lung injury in experimental gram-negative bacterial pneumonia. *J Immunol*. 2005;174(6):3676–85.
4. Doyle JF, Schortgen F. Should we treat pyrexia? And how do we do it? *Crit Care*. 2016;20(1):303.
5. Chiappini E, Venturini E, Remaschi G, Principi N, Longhi R, Tovo PA, Becherucci P, Bonsignori F, Esposito S, Festini F, et al. 2016 update of the Italian Pediatric Society Guidelines for Management of Fever in Children. *J Pediatr*. 2017;180:177–183e171.
6. Clericetti CM, Milani GP, Bianchetti MG, Simonetti GD, Fossali EF, Balestra AM, Bozzini MA, Agostoni C, Lava SAG. Systematic review finds that fever phobia is a worldwide issue among caregivers and healthcare providers. *Acta Paediatr*. 2019;108(8):1393–7.
7. Gaffney GR, Bereznicki LR, Bereznicki BJ. Knowledge, beliefs and management of childhood fever among nurses and other health professionals: a cross-sectional survey. *Nurse Educ Today*. 2021;97:104731.
8. Gehri M, Guignard E, Djahnine SR, Cotting JQ, Yersin C, Di Paolo ER, Krahenbuhl JD, Pannatier A. When fever, paracetamol? Theory and practice in a paediatric outpatient clinic. *Pharm World Sci*. 2005;27(3):254–7.
9. Souza Mvd, Damião EBC, Buchhorn SMM, Rossato LM. Non-pharmacological fever and hyperthermia management in children: an integrative review. *Acta Paulista de Enfermagem* 2021;34.
10. Vicens-Blanes F, Miró-Bonet R, Molina-Mula J. Analysis of Nurses' and Physicians' Attitudes, Knowledge, and Perceptions toward Fever in Children: A Systematic Review with Meta-Analysis. *Int J Environ Res Public Health* 2021, 18(23).
11. Wrotek S, LeGrand EK, Dzialuk A, Alcock J. Let fever do its job: the meaning of fever in the pandemic era. *Evol Med Public Health*. 2021;9(1):26–35.
12. Watts R, Robertson J, Thomas G. Nursing management of fever in children: a systematic review. *Int J Nurs Pract*. 2003;9(1):1–8.
13. El Hussein MT, Cuncannon A. Nursing students' transfer of learning from simulated clinical experiences into clinical practice: a scoping review. *Nurse Educ Today*. 2022;116:105449.
14. Patelarou AE, Mechili EA, Ruzafa-Martinez M, Dolezel J, Gotlib J, Skela-Savič B, Ramos-Morcillo AJ, Finotto S, Jarosova D, Smodiš M et al. Educational Interventions for Teaching Evidence-Based Practice to Undergraduate Nursing Students: A Scoping Review. *Int J Environ Res Public Health* 2020, 17(17).
15. Schmitt BD. Fever phobia: misconceptions of parents about fevers. *Am J Dis Child*. 1980;134(2):176–81.
16. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Res Psychol*. 2006;3(2):77–101.
17. Braun V, Clarke V. What can "thematic analysis" offer health and wellbeing researchers? *Int J Qual Stud Health Well-being*. 2014;9:26152.
18. Milani GP, Corsello A, Fadda M, Falvo I, Gianni ML, Marseglia GL, Cuppari C, Bruzzese E, Bianchetti MG, Schulz PJ et al. Perception, knowledge and attitude towards childhood fever: A survey among final-year medical students. *Br J Clin Pharmacol* 2022.
19. Martins M, Abecasis F. Healthcare professionals approach paediatric fever in significantly different ways and fever phobia is not just limited to parents. *Acta Paediatr*. 2016;105(7):829–33.
20. Greensmith L. Nurses' knowledge of and attitudes towards fever and fever management in one Irish children's hospital. *J Child Health Care*. 2013;17(3):305–16.
21. Poirier MP, Davis PH, Gonzalez-del Rey JA, Monroe KW. Pediatric emergency department nurses' perspectives on fever in children. *Pediatr Emerg Care*. 2000;16(1):9–12.
22. Edwards HE, Courtney MD, Wilson JE, Monaghan SJ, Walsh AM. Fever management practises: what pediatric nurses say. *Nurs Health Sci*. 2001;3(3):119–30.
23. Purssell E, Collin J. Fever phobia: the impact of time and mortality—a systematic review and meta-analysis. *Int J Nurs Stud*. 2016;56:81–9.
24. Cusson RM, Meehan C, Bourgault A, Kelley T. Educating the next generation of nurses to be innovators and change agents. *J Prof Nurs*. 2020;36(2):13–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.