

Case report

Open Access

Feasibility of using quadriceps-strengthening exercise to improve pain and sleep in a severely demented elder with osteoarthritis – a case report

Pao-Feng Tsai*¹, Kathy Richards^{1,2} and Richard FitzRandolph³

Address: ¹College of Nursing, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA, ²Central Arkansas Veterans Healthcare System, Little Rock, Arkansas, USA and ³Department of Radiology, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA

E-mail: Pao-Feng Tsai* - Tsai-paofeng@uams.edu; Kathy Richards - RichardsKathyC@uams.edu; Richard FitzRandolph - FitzrandolphRichard@uams.edu

*Corresponding author

Published: 2 October 2002

Received: 30 April 2002

BMC Nursing 2002, 1:1

Accepted: 2 October 2002

This article is available from: <http://www.biomedcentral.com/1472-6955/1/1>

© 2002 Tsai et al; licensee BioMed Central Ltd. This article is published in Open Access: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Osteoarthritis (OA) of the knee, which is prevalent among older adults in nursing homes, causes significant pain and suffering, including disturbance of nocturnal sleep. One nonpharmacologic treatment option is quadriceps-strengthening exercise, however, the feasibility of such a treatment for reducing pain from OA in severely demented elders has not been studied. This report describes our test of the feasibility of such an exercise program, together with its effects on pain and sleep, in a severely demented nursing home resident.

Case presentation: The subject was an elderly man with severe cognitive impairment (Mini-Mental Status Exam score 4) and knee OA (Kellgren-Lawrence radiographic grade 4). He was enrolled in a 5-week, 10-session standardized progressive-resistance training program to strengthen the quadriceps, and completed all sessions. Pain was assessed with the Western Ontario and MacMaster OA Index (WOMAC) pain subscale, and sleep was assessed by actigraphy.

The patient was able to perform the exercises, with a revision to the protocol. However, the WOMAC OA pain subscale proved inadequate for measuring pain in a patient with low cognitive functioning, and therefore the effects on pain were inconclusive. Although his sleep improved after the intervention, the influence of his medications and the amount of daytime sleep on his nighttime sleep need to be considered.

Conclusions: A quadriceps-strengthening exercise program for treating OA of the knee is feasible in severely demented elders, although a better outcome measure is needed for pain.

Background

Osteoarthritis (OA) is a highly prevalent, disabling condition afflicting 16 million older adults in the United States [1]. Nearly 70% of the elderly population show radiographic evidence of OA [2], and the prevalence of OA in-

creases with age [3]. Chronic pain and suffering from OA account for \$15.5 billion annually (in 1994 dollars) in health care expenditures [4]; and among all the potentially painful disorders in older persons, OA accounts for the greatest proportion of pain complaints [5]. Pain is an im-

portant predictor of functional limitation in persons with OA of the knee [6].

Some 45% to 65% of elderly nursing home residents suffer from OA [7,8]. The knee is one of the most commonly affected sites [9]. Undertreatment of pain in nursing home residents has serious consequences for the quality of their sleep [10], and the combination of unrelieved pain and sleep disturbance further exacerbates existing cognitive impairment [11] and leads to depression [12] and disruptive behavior [13], while also increasing the considerable burden and costs of caring for elders.

Treatment options for frail nursing home residents with OA are limited. Nursing home residents tend to be in poorer overall condition than their counterparts dwelling in the community, and pharmacologic intervention tends to exacerbate their already frail physical condition. An intervention that is easy to administer and has few side effects is thus needed for nursing home residents. One treatment option is an exercise program.

Recently, attention has focused on the quadriceps in the treatment of knee OA pain. The quadriceps mechanism is of key importance for walking, standing, and using stairs, and weakness in this muscle may cause impaired function. In addition, quadriceps weakness is a primary risk factor for progression of joint damage in knee OA and knee pain [9,14]. Slemenda et al. [15] found that less quadriceps strength predicted both radiographic and symptomatic knee OA. The odds ratio for the presence of OA per 10-lb-ft loss of strength was 0.8 for radiographic OA and it was 7.1 for symptomatic OA, indicating that persons with symptomatic OA had weaker quadriceps than those with asymptomatic OA. Further, O'Reilly et al. found that quadriceps weakness was strongly associated with pain in 600 community-dwelling individuals ages 40–79 with knee OA [9]. Subjects with knee pain had less voluntary quadriceps strength than those without pain ($t = 3.90$, $p < .01$). Quadriceps strength (Odds ratio = 18.8 for muscle strength ≤ 10 kgF) and radiographic change (Odds ratio = 4.1 for radiographic score ≥ 4) are thus independently associated with knee pain.

Exercise has been found to be an effective and well-tolerated treatment for knee OA [16,17]. Examining the effect of quadriceps exercise on knee OA in 113 subjects ages 50–80, Maurer et al. [16] found that both exercise and an educational intervention effected an overall improvement ($p < .05$). Patients in the exercise group, however, had a greater decrease in pain than those receiving an educational intervention ($p < .01$ for pain change; $p < .05$ for stair-associated pain at week 8). Similarly, Ettinger et al. [17], working with 365 subjects age 60 or older, reported

an 8% lower pain score for the quadriceps-exercise group than the educational-intervention group ($p < .05$).

Exercise in general is beneficial for nursing home residents with OA pain and disability. However, the feasibility of using a quadriceps-strengthening program for severely demented nursing home residents has not been tested. This case report describes our experience with a severely demented nursing home resident, the effects of quadriceps strengthening on his pain and sleep, and the implications of our experience for such an exercise program with this population.

Case presentation

Mr. T, a pleasant 80-year-old veteran, was chosen to participate in the exercise program. His medical diagnosis included dementia with delusional features, osteoarthritis, depression, anemia, gout, cataracts, and orthostatic hypotension. He was English-speaking, had a Mini-Mental Status Exam score of 4, had radiographic evidence of knee OA (Kellgren-Lawrence grade of 4 in both knees), and was ambulatory without assistance.

He was identified by nursing staff as having knee pain. Both he and his daughter provided consent for him to participate in the program. We collected information about his pain using the Western Ontario and MacMaster OA Index (WOMAC) pain subscale, and minutes of nighttime sleep were measured by actigraphy. Pain assessment occurred between 3:00 pm and 5:00 pm on days 1, 3, and 5 of baseline and days 16, 18, 20, 31, 33, and 35 of the intervention. Sleep assessment occurred on days 1–5 of baseline and days 16–20 and 31–35 of the intervention. These assessments were performed by the first author and a research assistant who was enrolled in a master's-level nurse practitioner program at the time of the study. Use of analgesic medications and medications that might affect sleep were recorded throughout the study period.

The 5-week quadriceps-strengthening exercise program involved standardized progressive-resistance training. After 5 minutes of warm-up, Mr. T performed knee extensions using Keiser knee-extension equipment. The exercise protocol included 10 sessions that required the elder to lift and lower the training loads separately for each leg to equalize the relative training stimulus for each limb. Mr. T performed three sets of eight repetitions on two nonsequential days per week for 5 weeks. His training loads began at 50% of his predetermined one-repetition maximum (1RM) and should gradually progress to heavier loads, which were decided based on Mr. T's tolerance of training loads and whether he showed any pain behaviors, such as refusing to exercise, guarding or rubbing the knee, showing nonverbal vocalizations (sighs, gasps, moans, groans, and cries) and facial grimacing or wincing,




Figure 1
The outcomes and use of medications throughout the project

and expressing vocal complaints (in words expressing discomfort or pain). All sessions were conducted with supervision to ensure safety and proper technique. A cool-down period of 5 minutes followed each exercise session. Mr. T completed all 10 quadriceps-strengthening exercise sessions.

At the pretest, his 1RM was 45 lb for both legs, which is similar to the average for osteoarthritic male elders [18] and is weaker than elders without OA [19]. We started with an average of 20 lb of resistance at the beginning of the program and maintained it throughout the intervention. The time required to complete a set of exercises varied depending on Mr. T's cognitive functioning on that day. At the beginning of the exercise intervention, it took an average of 3 to 4 minutes to complete a set because we needed to coach or help the confused elder do the exer-

cise. By the end of the exercise intervention, it took less than 1 minute for him to perform a set of exercises. Although we tried not to have a rest period between sets, we had no control over this. It depended on the cognitive functioning of Mr. T on that specific day.

After completing 10 sessions of exercises, his WOMAC pain score dropped from 3 at the pretest to 0, indicating decreased pain. His night-time sleep also improved, increasing from 271 minutes at the pretest to 474 minutes after the intervention. In addition, his Mini-Mental Status Exam score increased from 4 to 7. He remained ambulatory without assistance. His nutrition and appetite remained unchanged throughout the study period. The amount of anticonvulsant (valproic acid) that Mr. T was given increased from 350 mg to 500 mg during the last week of the intervention. In addition, his intake of anti-

psychotic (risperidone) increased from 0.65 mg to 1 mg after the second week of the intervention, while the amount of analgesic remained almost the same. Changes in his pain score, sleep, and use of medications are shown in Figure 1.

Because of his low cognitive functioning, during the first 2 weeks of the intervention, Mr. T could not understand the command for him to do each step of the exercise. We had to use our hands to nudge or touch his leg to give him a hint, and on occasion we lifted and lowered his legs for him. By the third week of the intervention, however, we saw a learning effect: Mr. T recognized the name of the machine (Keiser), pronouncing it every time we took him to the exercise laboratory, and he was able to sit on the machine by himself and perform the exercise almost without help. Throughout the exercise program, however, he could not exercise each leg separately.

The validity of self-reports of pain by elders with low cognitive functioning, such as Mr. T, is questionable. We had to keep asking the same questions over and over, and it was difficult to keep him focused on them. In addition, we had to ask the pain questions immediately after he performed each activity (walking, sitting, lying down, standing) because he could not remember what had happened more than a few minutes. When we asked the same questions on the WOMAC pain scale 5 minutes apart, he could not repeat the answers, which indicates poor test-retest reliability and questionable validity with the cognitively impaired.

Since the WOMAC pain scale is inadequate for elders with low cognitive functioning, alternative pain measures are needed. A possible alternative is assessment of activity level. We noted that Mr. T was usually very active and had good mobility, but when he suffered from pain in the lower extremities, he avoided activities (e.g., standing up, walking) or refused to do exercises. On day 2 of the intervention, we saw him sit in a chair, which was unusual; he then refused to stand up and pointed to his knee, saying, "It hurts." We stopped the intervention for that day and did it another day. Although Mr. T was able to communicate his pain on this occasion, often he could not do so. However, an unusual activity level in itself may be an indicator of pain in demented elders with OA of the lower extremities.

Mr. T tolerated the actigraph well. He was curious about the wristwatch-like device secured to his wrist with a strip, touching it and playing with it. He kept the actigraph on his wrist for the required measurement period. Although his sleep improved, his medications (valproic acid and risperidone) may have influenced his sleep (Figure 1). Future study needs to control for the use of medications, if

possible. We also suspected that the total minutes of daytime sleep may have been associated with sleep time during the night (Figure 1). During the last week of the intervention, Mr. T slept about 127 minutes during the day, which may have contributed to lack of sleep during the night (192 minutes). On day 3 of the last week of intervention, Mr. T slept only 28 minutes during the day, and this may have contributed to the greater amount of nighttime sleep.

Conclusions

Even with severe cognitive impairment, Mr. T was able to perform the quadriceps-strengthening exercises. The exercise program, however, needs to be adapted to an individual's level of cognitive functioning (e.g., using both legs simultaneously in the exercise instead of each leg alternately). The pain subscale of the WOMAC OA Index is inadequate for measuring pain in elders with low cognitive functioning, who have difficulty communicating pain. In addition, the reliability and validity of self-reports of pain by these elders are questionable. It is necessary to find an alternative means of assessing pain in this population. Although Mr. T's sleep improved after the intervention, the influence of medications and the amount of daytime sleep need to be considered. The actigraph appeared to be well tolerated and is an appropriate device for measuring the sleep of demented elders.

In conclusion, a quadriceps-strengthening exercise program for treating OA of the knee is feasible with severely demented elders. However, such an exercise program is labor intensive and an expensive intervention because severely demented elders will need to be closely supervised by the staff. With the shortage of nursing staffs, many nursing homes may not have the capacity to implement the entire exercise program. We learned from this project that severely demented elders are able to learn, able to follow directions and able to participate in an exercise program. Therefore, we suggest modifying the exercise program to make it a group activity and teach demented elders with OA of the knee to perform leg-extension exercises without an exercise machine. Further investigation will be needed to determine the effects of a modified exercise program, and a better measure of pain is needed.

List of abbreviations used

Osteoarthritis: OA

Western Ontario and MacMaster Osteoarthritis Index: WOMAC

Predetermined one-repetition maximum: 1RM

Declaration of competing interests

None of the authors receives reimbursements, fees, funding, or salary from an organisation or held any stocks or shares in an organisation that may in any way gain or lose financially from the publication of this paper in the past five years. None of the authors has any nonfinancial competing interests they would like to declare in relation to this paper.

Authors' contributions

Author 1 PT developed the research proposal, carried out the actual exercise program, and participated in the sequence alignment of the manuscript. Author 2 KR participated in developing the research proposal and interpreting the sleep data. Author 3 RF read the radiographs and performed the diagnosis of OA.

All authors read and approved the final manuscript.

Acknowledgements

The first author was supported by the Intramural Grant Program and the Hartford Center of Geriatric Nursing Excellence, College of Nursing, University of Arkansas for Medical Sciences while this research was being conducted. We thank Ms. Pamela Avaltroni and Melissa Grubbs for their assistance during the data collection. We also thank Ms. Elizabeth Tornquist and Mr. William Gabello for editorial assistance during the preparation of this manuscript. Finally, we thank Mr. T and his family for participating in this project. Written consent was obtained from the patient and his family for publication of the patient's details.

References

1. American College of Rheumatology: **The American College of Rheumatology Clinical Guidelines**. Atlanta, Georgia: American College of Rheumatology 1997
2. Lawrence RC, Hochberg MC, Kelsey JL, et al: **Estimates of the prevalence of selected arthritic and musculoskeletal diseases in the United States**. *Journal of Rheumatology* 1989, **16(4)**:427-41
3. Hamerman D: **Clinical implications of osteoarthritis and aging**. *Annals of the Rheumatic Diseases* 1995, **54(2)**:82-5
4. Yelin E: **The economics of osteoarthritis**. In: *Osteoarthritis* (Edited by: Brandt K, Doherty M, Lohmander LS) New York: Oxford University Press 1998, 23-30
5. Sternbach RA: **Survey of Pain in the United States: The Nuprin Pain Report**. *Clin J Pain* 1986, **2**:49-53
6. Hochberg MC, Lawrence RC, Everett DF, Cornoni-Huntley J: **Epidemiologic associations of pain in osteoarthritis of the knee: data from the National Health and Nutrition Examination Survey and the National Health and Nutrition Examination I Epidemiologic Follow-up Survey**. *Seminars in Arthritis & Rheumatism* 1989, **8(4 Suppl 2)**:4-9
7. Marzinski LR: **The tragedy of dementia: clinically assessing pain in the confused nonverbal elderly**. *Journal of Gerontological Nursing* 1991, **17(6)**:25-8
8. Ferrell BA, Ferrell BR, Osterweil D: **Pain in the nursing home**. *Journal of the American Geriatrics Society* 1990, **38(4)**:409-14
9. O'Reilly SC, Jones A, Muir KR, Doherty M: **Quadriceps weakness in knee osteoarthritis: the effect on pain and disability**. *Annals of the Rheumatic Diseases* 1998, **57(10)**:588-94
10. Ross MM, Crook J: **Elderly recipients of home nursing services: pain, disability and functional competence**. *Journal of Advanced Nursing* 1998, **27(6)**:1117-26
11. Duggleby W, Lander J: **Cognitive status and postoperative pain: older adults**. *Journal of Pain & Symptom Management* 1994, **9(1)**:19-27
12. Cohen-Mansfield J, Marx MS: **Pain and depression in the nursing home: corroborating results**. *Journal of Gerontology* 1993, **48(2)**:96-7
13. Ryden MB, Bossenmaier M, McLachlan C: **Aggressive behavior in cognitively impaired nursing home residents**. *Research in Nursing & Health* 1991, **14(2)**:87-95
14. Slemenda C, Brandt KD, Heilman DK, et al: **Quadriceps weakness and osteoarthritis of the knee**. *Annals of Internal Medicine* 1997, **127(2)**:97-104
15. Slemenda C, Heilman DK, Brandt KD, et al: **Reduced quadriceps strength relative to body weight: a risk factor for knee osteoarthritis in women?** *Arthritis & Rheumatism* 1998, **41(11)**:1951-9
16. Maurer BT, Stern AG, Kinossian B, Cook KD, Schumacher HR Jr: **Osteoarthritis of the knee: isokinetic quadriceps exercise versus an educational intervention**. *Archives of Physical Medicine & Rehabilitation* 1999, **80(10)**:1293-9
17. Ettinger WH Jr, Burns R, Messier SP, et al: **A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST)** *JAMA* 1997, **277(1)**:25-31
18. Brandt KD, Heilman DK, Slemenda C, et al: **Quadriceps strength in women with radiographically progressive osteoarthritis of the knee and those with stable radiographic changes**. *J Rheumatol* 1999, **26(11)**:2431-7
19. Trappe S, Williamson D, Godard M: **Maintenance of whole muscle strength and size following resistance training in older men**. *J Gerontol A Biol Sci Med Sci* 2002, **57(4)**:B138-43

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1472-6955/1/1/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



BioMedcentral.com

Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com